

THE EVIDENCE BEHIND **CUES** AN INTERVENTION TO ADDRESS INTIMATE PARTNER VIOLENCE IN HEALTH SETTINGS

With the CUES intervention, healthcare professionals talk with all patients about how relationships can affect health and how to get support. Unlike traditional screening methods, CUES does not rely on a disclosure of abuse in order to provide a patient with information and resources they might need. By using CUES, providers can ensure that patients are aware of available survivor support services and have information to pass to friends and family who might need it. In order to adopt a team-based approach to supporting survivors, this approach relies on fostering of strong partnerships between domestic violence agencies and health settings.

CUES stands for Confidentiality, Universal Education + Empowerment, and Support

C: Confidentiality

- ▶ Know your state's reporting requirements and share any limits of confidentiality with your patients.
- ▶ Ensure that you can bring up relationships and violence safely by seeing patients alone for at least part of the in person or telehealth visit.

UE: Universal Education + Empowerment

- ▶ In-Person Visits: Share two safety cards with each patient to start the conversation about relationships and how they affect health. Make sure patients know that you're a safe person for them to talk to. Encourage patients to share a safety card with a friend or family member.
- ▶ Telehealth visits: Normalize with all patients that stress from relationship struggles or previous trauma can affect health. Ask if you can send them a link to resources should they need them for themselves or a friend or family member.



Safety cards are available for different settings, communities, and languages.

S: Support

- ▶ Though disclosure of violence is not the goal, it will happen – know how to support someone who shares that they have experienced abuse.
- ▶ Offer health promotion strategies and a care plan that takes surviving abuse into consideration.
- ▶ Make a warm referral to your local domestic/sexual violence partner agency or national hotlines (on the back of all safety cards).

Futures Without Violence National Health Resource Center on Domestic Violence

The building blocks of the CUES intervention have been evaluated in reproductive and adolescent health settings by Elizabeth Miller, MD, PhD. The development of this intervention was based on research done over the last fifteen years about addressing intimate partner violence in healthcare.

What Survivors of IPV Want from Health Professionals

Autonomy Survivors want to be able to make their own decisions when it comes to health care.¹⁻⁵ This includes choosing their own providers, disclosing in their own time, and having multiple pathways for potential interventions.^{1,4,5} Additionally, health professionals should always explain procedures thoroughly and allow patients to ask questions or signal discomfort.¹ Intimate partner violence can strip people of control—provider awareness can serve as a powerful tool to combat this.⁵

Empathy & Compassion People who have been exposed to intimate partner violence want providers to validate their experiences.^{1,2,4} Providers should strive to be nonjudgmental and supportive, regardless of how survivors of intimate partner violence respond to interventions.^{4,5} A judgmental, intrusive or controlling provider may recall trauma for survivors.¹

Informed Providers Providers with knowledge about intimate partner violence make a significant difference in the experiences of patients. Survivors cite the value of having health professionals who understand the depth and complexity of IPV. This includes the impact of trauma on health, the long-term nature of this violence, and its intersection with accessing a variety of other needs (employment, childcare, transportation, etc.)^{1,4,5}

Limitations of Screening for IPV without Universal Education

Low Disclosure Rates Disclosure rates among patients screened for intimate partner violence in health care settings are not the same as known prevalence of intimate partner violence from research studies. While 1 in 4 women and 1 in 9 men are estimated to be survivors of intimate partner violence in the general population, disclosure rates in clinical settings range from 1%-14% with screening, typically hovering around 7%.⁶⁻¹⁰ Among known survivors, only 21.1% of women and 5.6% of men report disclosing to a healthcare professional.⁶

Non-Differential Outcomes for Screening Alone Outcomes for survivors are comparable between screened and non-screened patients in the absence of universal education or warm referrals.¹¹ No significant differences emerge in quality of life, number of emergency room visits, or number of hospitalizations.¹² Beyond this, no significant differences have been found in referral behaviors.⁹

Importance of Confidentiality and Patients' Reasons for Non-Disclosure

Patient Concerns Patients who have experienced IPV are concerned about the privacy of their data, particularly with electronic medical records.¹³ A perceived lack of privacy is a reason for non-disclosure by survivors.¹³⁻¹⁶ Assured confidentiality of providers, as well as a clear explanation of how patient information will be used, is recommended.¹³

Reasons for Non-Disclosure Amongst patients with a history of IPV, reasons for non-disclosure include fear of judgment, emotional distress, fear of not receiving adequate support from healthcare professionals, concern about mandated reporting procedures, religious beliefs, language barriers, and having children in the home.¹⁴⁻¹⁶

Value of Universal Education & Empowerment

Universal Education and Empowerment The National Sexual Violence Resource Center encourages a trauma-informed framework that focuses on being sensitive to potential trauma in survivors, rather than responding solely to disclosure.¹⁷ Comprehensive approaches including education, assessment, intervention, and prevention are recommended to empower and uplift survivors.^{18,19}

Outcomes for Patients In an intervention centered around reducing unintended pregnancies in women who experience intimate partner violence, utilization of a universal education model was associated with a 71% reduction in pregnancy pressure, and women being 60% more likely to end a relationship because it felt unsafe.^{18,20} Additional universal education models have demonstrated a threefold increase in disclosure amongst youth who experienced relationship abuse, as well as decreased self-reported isolation and reduced reproductive coercion in women with high baselines of RC.^{21,22} Patients exposed to these interventions have positive feedback, and report them to be more helpful than comparable resources.²³

Outcomes for Providers Providers exposed to a universal education and empowerment curriculum report improved confidence in discussing IPV, as well as increased awareness about the complexity of issues faced by survivors.²³ These outcomes are cited as reasons providers are less likely to screen.^{7, 24}

Peer-to-Peer Education Patients who receive universal education do not only directly benefit, but also share the information with their peers. Studies have demonstrated that participants who received UE were almost twice as likely to share the number for the domestic violence hotline with someone.²² Additionally, patients who receive palm-sized cards have cited the empowerment they feel when sharing concrete resources with their peers.²⁰

Cross-Sector Partnerships & Survivor Health and Safety

Coordinating Care Systems approaches, rooted in strong cross-sector networks, show important outcomes for women who have experienced IPV. Community partnerships create an easier “warm referral” process, allowing providers to facilitate the connection between survivors and resources. These warm referrals increase the likelihood of survivors utilizing interventions.²⁵

Collaborative Learning Network models and cross-sectoral curricula demonstrate an improved knowledge base for all participants. Collaborative groups successfully challenge misconceptions around intimate partner violence and built meaningful relationships.²⁶⁻²⁸ Culturally competent care is also improved through these collaborations.²⁷ These networks foster easier referral processes and stronger understandings of the field.²⁶⁻²⁸

National Health Resource Center on Domestic Violence at Futures Without Violence

Learn more: ipvhealth.org

For community health centers: ipvhealthpartners.org

Order [safety cards](#) and more: store.futureswithoutviolence.org

Email: health@futureswithoutviolence.org



References

1. Reeves EA, Humphreys JC. Describing the healthcare experiences and strategies of women survivors of violence. *J Clin Nurs*. 2018;27(5-6):1170-1182. doi:10.1111/jocn.14152
2. Munro-Kramer ML, Dulin AC, Gaither C. What survivors want: Understanding the needs of sexual assault survivors. *J Am Coll Health*. 2017;65(5):297-305. doi:10.1080/07448481.2017.1312409
3. Kulkarni SJ, Bell H, Rhodes DM. Back to basics: essential qualities of services for survivors of intimate partner violence. *Violence Against Women*. 2012;18(1):85-101. doi:10.1177/1077801212437137
4. Feder GS, Hutson M, Ramsay J, Taket AR. Women exposed to intimate partner violence: expectations and experiences when they encounter health care professionals: a meta-analysis of qualitative studies. *Arch Intern Med*. 2006;166(1):22-37. doi:10.1001/archinte.166.1.22
5. Chang JC, Cluss PA, Ranieri L, et al. Health care interventions for intimate partner violence: what women want. *Womens Health Issues*. 2005;15(1):21-30. doi:10.1016/j.whi.2004.08.007
6. Black MC, Basile KC, Breiding MJ, et al. The national intimate partner and sexual violence survey: 2010 summary report. Atlanta, GA Natl Cent Inj Prev Control Centers Dis Control Prev. 2011;19:39-40.
7. Ahmad I, Ali PA, Rehman S, Talpur A, Dhingra K. Intimate partner violence screening in emergency department: a rapid review of the literature. *J Clin Nurs*. 2017;26(21-22):3271-3285. doi:10.1111/jocn.13706
8. Hussain N, Sprague S, Madden K, Hussain FN, Pindiprolu B, Bhandari M. A comparison of the types of screening tool administration methods used for the detection of intimate partner violence: a systematic review and meta-analysis. *Trauma Violence Abuse*. 2015;16(1):60-69. doi:10.1177/1524838013515759
9. Taft A, O'Doherty L, Hegarty K, Ramsay J, Davidson L, Feder G. Screening women for intimate partner violence in healthcare settings. *Cochrane database Syst Rev*. 2013;(4):CD007007. doi:10.1002/14651858.CD007007.pub2
10. Feder G, Davies RA, Baird K, et al. Identification and Referral to Improve Safety (IRIS) of women experiencing domestic violence with a primary care training and support programme: a cluster randomised controlled trial. *Lancet (London, England)*. 2011;378(9805):1788-1795. doi:10.1016/S0140-6736(11)61179-3
11. Wathen CN, MacMillan HL. Health care's response to women exposed to partner violence: moving beyond universal screening. *JAMA*. 2012;308(7):712-713. doi:10.1001/jama.2012.9913
12. Klevens J, Kee R, Trick W, et al. Effect of screening for partner violence on women's quality of life: a randomized controlled trial. *JAMA*. 2012;308(7):681-689. doi:10.1001/jama.2012.6434
13. Dichter ME, Wagner C, Goldberg EB, Iverson KM. Intimate Partner Violence Detection and Care in the Veterans Health Administration: Patient and Provider Perspectives. *Womens Health Issues*. 2015;25(5):555-560. doi:10.1016/j.whi.2015.06.006
14. Nelson HD, Bougatsos C, Blazina I. Screening women for intimate partner violence: a systematic review to update the US Preventive Services Task Force recommendation. *Ann Intern Med*. 2012;156(11):796-808.
15. Othman S, Goddard C, Piterman L. Victims' barriers to discussing domestic violence in clinical consultations: a qualitative enquiry. *J Interpers Violence*. 2014;29(8):1497-1513. doi:10.1177/0886260513507136

16. Montalvo-Liendo N. Cross-cultural factors in disclosure of intimate partner violence: an integrated review. *J Adv Nurs*. 2009;65(1):20–34. doi:10.1111/j.1365-2648.2008.04850.x
17. The National Sexual Assault Coalition Resource Sharing Project and National Sexual Violence Resource Center. Building Cultures of Care: A Guide for Sexual Assault Services Programs.; 2013. https://www.nsvrc.org/sites/default/files/2017-10/publications_nsvrc_building-cultures-of-care.pdf.
18. Decker MR, Frattaroli S, McCaw B, et al. Transforming the healthcare response to intimate partner violence and taking best practices to scale. *J Womens Health (Larchmt)*. 2012;21(12):1222–1229. doi:10.1089/jwh.2012.4058
19. Ghandour RM, Campbell JC, Lloyd J. Screening and counseling for Intimate Partner Violence: a vision for the future. *J Womens Health (Larchmt)*. 2015;24(1):57–61. doi:10.1089/jwh.2014.4885
20. Miller E, Decker MR, McCauley HL, et al. A family planning clinic partner violence intervention to reduce risk associated with reproductive coercion. *Contraception*. 2011;83(3):274–280. doi:10.1016/j.contraception.2010.07.013
21. Miller E, Goldstein S, McCauley HL, et al. A school health center intervention for abusive adolescent relationships: a cluster RCT. *Pediatrics*. 2015;135(1):76–85. doi:10.1542/peds.2014-2471
22. Miller E, Tancredi DJ, Decker MR, et al. A family planning clinic-based intervention to address reproductive coercion: a cluster randomized controlled trial. *Contraception*. 2016;94(1):58–67. doi:10.1016/j.contraception.2016.02.009
23. Miller E, McCauley HL, Decker MR, et al. Implementation of a Family Planning Clinic-Based Partner Violence and Reproductive Coercion Intervention: Provider and Patient Perspectives. *Perspect Sex Reprod Health*. 2017;49(2):85–93. doi:10.1363/psrh.12021
24. Williams JR, Halstead V, Salani D, Koermer N. An exploration of screening protocols for intimate partner violence in healthcare facilities: a qualitative study. *J Clin Nurs*. 2017;26(15-16):2192–2201. doi:10.1111/jocn.13353
25. Miller E, McCaw B, Humphreys BL, Mitchell C. Integrating intimate partner violence assessment and intervention into healthcare in the United States: a systems approach. *J Womens Health (Larchmt)*. 2015;24(1):92–99. doi:10.1089/jwh.2014.4870
26. Whitaker DJ, Baker CK, Pratt C, et al. A network model for providing culturally competent services for intimate partner violence and sexual violence. *Violence Against Women*. 2007;13(2):190–209. doi:10.1177/1077801206296984
27. Mason R, Wolf M, O'Rinn S, Ene G. Making connections across silos: intimate partner violence, mental health, and substance use. *BMC Womens Health*. 2017;17(1):29. doi:10.1186/s12905-017-0372-4
28. Laing L, Irwin J, Toivonen C. Across the divide: using research to enhance collaboration between mental health and domestic violence services. *Aust Soc Work*. 2012;65(1):120–135.
29. Chung EK, Siegel BS, Garg A, et al. Screening for social determinants of health among children and families living in poverty: a guide for clinicians. *Curr Probl Pediatr Adolesc Health Care*. 2016;46(5):135–153.
30. Garg A, Boynton-Jarrett R, Dworkin PH. Avoiding the unintended consequences of screening for social determinants of health. *Jama*. 2016;316(8):813–814.
31. Billioux A, Verlander K, Anthony S, Alley D. Standardized screening for health-related social needs in clinical settings: The accountable health communities screening tool. *NAM Perspect*. 2017.

32. United States Department of Agriculture Food and Nutrition Service. Guide to Measuring Household Food Security (Revised 2000). <https://www.fns.usda.gov/guide-measuring-household-food-security-revised-2000>. Published 2013.
33. Hager ER, Quigg AM, Black MM, et al. Development and validity of a 2-item screen to identify families at risk for food insecurity. *Pediatrics*. 2010;126(1):e26–e32.
34. Rosa TL, Ortolano SE, Dickin KL. Remembering food insecurity: Low-income parents' perspectives on childhood experiences and implications for measurement. *Appetite*. 2018;121:1–8.
35. Williams DR, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental health: Socio-economic status, stress and discrimination. *J Health Psychol*. 1997;2(3):335–351.
36. Williams DR, Gonzalez HM, Williams S, Mohammed SA, Moomal H, Stein DJ. Perceived discrimination, race and health in South Africa. *Soc Sci Med*. 2008;67(3):441–452.
37. McNeilly MD, Anderson NB, Armstead CA, et al. The perceived racism scale: a multidimensional assessment of the experience of white racism among African Americans. *Ethn Dis*. 1996.
38. Sandel M, Sheward R, de Cuba SE, et al. Unstable housing and caregiver and child health in renter families. *Pediatrics*. 2018;141(2):e20172199.
39. Byrne T, Fargo JD, Montgomery AE, Roberts CB, Culhane DP, Kane V. Screening for homelessness in the Veterans Health Administration: monitoring housing stability through repeat screening. *Public Health Rep*. 2015;130(6):684–692.
40. Syed ST, Gerber BS, Sharp LK. Traveling towards disease: transportation barriers to health care access. *J Community Health*. 2013;38(5):976–993. doi:10.1007/s10900-013-9681-1